

Review

Vascular Permeability Factor/Vascular Endothelial Growth Factor, Microvascular Hyperpermeability, and Angiogenesis

Harold F. Dvorak, Lawrence F. Brown,
Michael Detmar, and Ann M. Dvorak

*From the Departments of Pathology, Beth Israel Hospital
and Harvard Medical School, Boston, Massachusetts*

Vascular permeability factor (VPF), also known as vascular endothelial growth factor (VEGF) and vasculotropin, is a potent, multifunctional cytokine that exerts several important and possibly independent actions on vascular endothelium.¹ VPF/VEGF was originally discovered as a tumor-secreted protein that rendered venules and small veins hyperpermeable to circulating macromolecules.²⁻⁸ Subsequently it was learned that VPF/VEGF acts directly on cultured endothelial cells (ECs) to induce transient accumulation of cytoplasmic calcium,⁹ shape change,¹⁰ cell division,¹¹⁻¹⁹ and migration.²⁰ In addition, VPF/VEGF alters the pattern of EC gene expression^{21,22} and induces angiogenesis *in vivo*.^{11,23,24}

It is this last property, its capacity to induce angiogenesis, that has excited the greatest amount of current interest in VPF/VEGF. Angiogenesis is a complex process that involves EC division, selective degradation of vascular basement membranes and of surrounding extracellular matrix, and EC migration.²⁵ Directly or indirectly, VPF/VEGF has the ability to affect all of these activities. Moreover, both VPF/VEGF and its two specific EC receptors are commonly overexpressed at sites of new blood vessel growth. Taken together, VPF/VEGF has come to be recognized as an important angiogenesis factor.

Puzzling to some, however, has been the potent activity of VPF/VEGF as a vascular permeabilizing agent. Microvascular hyperpermeability has not generally been regarded as a component of angiogenesis. Therefore, it has seemed incongruous that the vascular permeabilizing effect of VPF/VEGF could be

related to its pro-angiogenic activity. However, as we shall recount, there is a large body of evidence indicating that microvascular hyperpermeability regularly and perhaps invariably precedes and/or accompanies angiogenesis as it occurs in a wide variety of pathological and physiological settings. Moreover, investigations into the consequences of microvascular hyperpermeability provide a rationale for its importance in vascular neogenesis.

In this review we briefly summarize the properties and functions of VPF/VEGF and its association with angiogenesis in tumors, in non-neoplastic pathologies such as wound healing and inflammation, and in certain physiological processes. In each of these settings there is good evidence that angiogenesis is accompanied by VPF/VEGF-mediated microvascular hyperpermeability. Taken together, the data indicate that microvascular hyperpermeability regularly accompanies and likely has a mechanistic role in the induction of angiogenesis.

VPF/VEGF Structure and Functions

VPF/VEGF is a highly conserved, disulfide-bonded dimeric glycoprotein of Mr 34 to 45 kd, which, upon reduction, loses all of its biological activity and separates into major bands of 17 to 22 kD.^{6-8,12,15,26-28} VPF/VEGF shares low but significant sequence homology with platelet-derived growth factor (PDGF) and closer homology with placenta growth factor.²⁹

Supported by USPHS NIH grants CA-50453 and CA-58845 from the National Cancer Institute, by the National Psoriasis Foundation, by the BIH Pathology Foundation, Inc., and under terms of a contract from the National Foundation for Cancer Research.

Accepted for publication February 13, 1995.

Address reprint requests to Harold F. Dvorak, MD, Dept. of Pathology, Beth Israel Hospital, Boston, MA 02215

Four different VPF/VEGF transcripts encoding polypeptides of 206, 189, 165, and 121 amino acids may be expressed by human cells.^{30,31} VPF/VEGF₁₆₅ is expressed predominantly in most instances. However, all isoforms arise from alternative splicing of a single gene whose coding region is divided among eight exons; typical of a secreted protein, the DNA structure predicts a hydrophobic leader sequence that is absent from the mature protein.³⁰ The several different VPF/VEGF isoforms apparently express identical biological activities. However, it appears that VPF/VEGF₁₂₁ and to a large extent VPF/VEGF₁₆₅ are secreted in soluble form, whereas the two larger isoforms remain cell-associated, perhaps because of their greater affinity for cell-surface proteoglycans.^{31,32} VPF/VEGF was originally purified on the basis of its affinity for heparin.^{4,8} Nonetheless, the affinity of VPF/VEGF₁₆₅ for heparin is substantially lower than that of other typical heparin-binding growth factors such as basic fibroblast growth factor⁴ and the smaller VPF/VEGF₁₂₁ isoform does not bind to heparin.³²

VPF/VEGF acts directly and selectively on vascular ECs by way of two Class III receptor tyrosine kinases, flt-1 and KDR, that are expressed predominantly, if not exclusively, on vascular endothelium. Several effects follow VPF/VEGF interaction with EC receptors. The earliest to be detected is an increase up to fourfold of cytoplasmic calcium ($[Ca^{2+}]_i$); $[Ca^{2+}]_i$ begins to increase within seconds after exposure of cultured ECs to subpicomolar concentrations of VPF/VEGF, making this measurement the most sensitive assay currently available for detecting VPF/VEGF activity.⁹ The effect of VPF/VEGF on EC $[Ca^{2+}]_i$ resembles that of other EC agonists such as thrombin and histamine and includes increased release of von Willebrand factor; however, as noted above, VPF/VEGF acts by way of distinct EC receptors, and its action is unaffected by inhibitors of thrombin and histamine.⁹ Like other agonists that increase $[Ca^{2+}]_i$, VPF/VEGF stimulates IP_3 accumulation and is thought to act through a phosphoinositide-specific phospholipase C.⁹

VPF/VEGF was originally discovered because of its ability to increase the permeability of microvessels, primarily postcapillary venules and small veins, to circulating macromolecules. It is one of the most potent vascular permeabilizing agents known, acting at concentrations below 1 nmol/L, and, as measured in the Miles assay, with a potency some 50,000 times that of histamine.^{1,2,4,10,33} VPF/VEGF acts to permeabilize a number of vascular beds, including those of the skin, peritoneal wall, mesentery, and diaphragm.^{2,4,34} The vascular permeabilizing effect occurs rapidly, be-

coming evident within several minutes of VPF/VEGF injection.^{2,4,34} Like its effects on EC $[Ca^{2+}]_i$, the vascular hyperpermeability resulting from a single injection of VPF/VEGF is transient and reversible, persisting for <30 minutes, and is not associated with detectable injury to ECs or to other microvascular components. The vascular permeabilizing action of VPF/VEGF is not blocked by antihistamines or by any of a variety of inhibitors of inflammation including those that inhibit platelet activating factor^{2,4,33} (D. Senger, unpublished data).

VPF/VEGF apparently increases microvascular permeability by enhancing the functional activity of a recently described organelle, the vesicular-vacuolar organelle (VVO).^{35,36} VVOs are grape-like clusters of uncoated vesicles and vacuoles deployed at intervals in the cytoplasm of ECs lining venules and small veins. The individual vesicles and vacuoles comprising VVOs are bounded by trilaminar unit membranes and are reminiscent of the 60 to 80-nm vesicles or caveolae that have been described in capillary endothelium. In contrast to caveolae, VVOs are elaborate structures, each of which, in a single section plane, comprises an average of 12 (up to as many as 20) individual vesicles or vacuoles. Commonly, VVOs span the entire thickness of EC cytoplasm from the luminal to the abluminal plasma membranes. The individual vesicles and vacuoles that make up VVOs interconnect with each other and with the EC plasma membranes by means of fenestrae that may be open or closed by diaphragms. When fenestrae are open, macromolecular tracers are able to pass between interconnecting vesicles and vacuoles, and their flow can be followed at sequential intervals across the EC cytoplasm from the vascular lumen to the abluminal basal lamina; thus, VVOs provide a pathway whereby plasma and plasma proteins may exit the circulation and enter the tissues.

In normal adult tissues, extravasation of circulating macromolecules from the microvasculature is quite limited but, to the extent that it occurs, takes place largely (eg, horseradish peroxidase) or entirely (eg, ferritin, dextrans) by way of VVOs.³⁵ In tumor vessels, or in normal skin following local injection of VPF/VEGF, VVO function is substantially upregulated (unpublished data). The mechanisms by which VPF/VEGF upregulates VVO function remain to be determined, but VPF/VEGF has been localized by immunocytochemistry to the abluminal plasma membrane and to VVOs of tumor vessel ECs.³⁶ Together these data suggest that VPF/VEGF may act to open fenestrae, thereby facilitating tracer extravasation. They also suggest that VVOs may correspond to the "large

pores" that physiologists have postulated in endothelial cells to account for the transendothelial transport of macromolecules.³⁷

In addition to increasing EC cytoplasmic $[Ca^{2+}]$, and permeability to macromolecules, VPF/VEGF causes ECs from several different sources to assume an elongated shape and stimulates their replication.¹⁰⁻¹⁹ VPF/VEGF also stimulates the migration of ECs,^{20,38} mouse monocytes,³⁹ and fetal bovine osteoblasts.⁴⁰ To our knowledge, these last are the only examples thus far in which VPF/VEGF exerts a direct biological effect on cells other than vascular endothelium. More generally, VPF/VEGF alters the pattern of EC gene activation, upregulating the expression of both plasminogen activators (PAs), uPA, and tPA, as well as the PA inhibitor PAI-1.²¹ VPF/VEGF also induces the expression of another EC protease, interstitial collagenase,²² and tissue factor.³⁹ Finally, and of particular relevance to this discussion, VPF/VEGF promotes angiogenesis in a variety of assay systems, both *in vivo* and *in vitro*.^{11,23,24}

Expression and Distribution of VPF/VEGF and its Receptors in Angiogenesis

VPF/VEGF Expression

VPF/VEGF is substantially overexpressed at both the mRNA and protein levels in a high percentage of malignant animal and human tumors,^{2-4,41-44} as well as in many transformed cell lines.⁵ Autochthonous human tumors that overexpress VPF/VEGF include a majority of carcinomas arising in the colon, stomach, pancreas, kidney, bladder, and breast, and glioblastomas; exceptions include lobular carcinomas of the breast and papillary kidney tumors that express little or no VPF/VEGF mRNA as detected by *in situ* hybridization.^{42,44} Benign tumors have been less carefully studied but benign adenomatous polyps arising in the colon do not express VPF/VEGF above levels found in normal colonic mucosa.⁴¹ However, VPF/VEGF mRNA is overexpressed by ductal carcinomas *in situ* of the breast before evidence of tumor invasion.⁴⁴

VPF/VEGF is also overexpressed prominently in certain important non-neoplastic pathological states that, like tumors, are characterized by angiogenesis. Thus, epidermal keratinocytes, as well as a subpopulation of macrophage-like cells, overexpress VPF/VEGF mRNA in healing cutaneous wounds,⁴⁵ psoriasis,⁴⁶ and delayed-type hypersensitivity.⁴⁷ Similarly, cardiac myocytes overexpress VPF/VEGF in

ischemic myocardium^{48,49} as do synovial lining cells in the highly vascularized pannus of rheumatoid arthritis.^{38,50}

Finally, VPF/VEGF is expressed at high levels by the placenta,⁵¹⁻⁵³ by many fetal tissues,^{54,55} and by a number of normal adult tissues undergoing physiological angiogenesis; eg, in the proliferating endometrium⁵⁶ and in the corpus luteum.^{57,58} VPF/VEGF is also expressed at low levels in a wide variety of normal adult human and animal tissues and at higher levels in a few select sites, namely, podocytes of the renal glomerulus, cardiac myocytes, prostatic epithelium and semen, and certain epithelial cells of the adrenal cortex and lung.^{53,59-61} High VPF/VEGF expression in these last sites is exceptional in that they are not characterized by new blood vessel formation.

Recent studies have sought to determine the mechanisms regulating VPF/VEGF expression. As noted above, constitutive overexpression of VPF/VEGF has been associated with the transformed phenotype and with vascularized, autochthonous, and transplantable tumors growing *in vivo*. VPF/VEGF is also overexpressed cyclically in certain processes involving neovascularization (menstrual cycle, corpus luteum formation, etc.) that are under hormonal control. In addition, certain cytokines and growth factors stimulate VPF/VEGF mRNA expression; apparently these effects are target cell specific in that transforming growth factor- α (TGF- α), and to a lesser extent epidermal growth factor, are stimulatory in keratinocytes,⁴⁶ whereas other cytokines such as TGF- β and PDGF upregulate VPF/VEGF expression in other types of cells.⁶²⁻⁶⁵ In all of these instances, VPF/VEGF overexpression is associated with angiogenesis.

Finally, tissue hypoxia has long been associated with angiogenesis, and there is good evidence that local oxygen concentrations regulate VPF/VEGF expression; in fact, the VPF/VEGF sequence shares elements with another oxygen-sensitive gene, erythropoietin.⁶⁶ In hypoxic tissues, VPF/VEGF mRNA expression is often increased above already high pre-existing levels; eg, in viable tumor cells situated immediately adjacent to zones of tumor necrosis, in hypoxic myocardium, etc.^{41-44,48,49,67} Moreover, in tumors with foci of relative hypoxia, VPF/VEGF mRNA may be expressed not only by malignant cells but also by stromal cells.^{41,42} These findings are consistent with those of *in vitro* experiments where limiting oxygen concentration provokes a number of cell types to overexpress VPF/VEGF mRNA.⁶⁶⁻⁶⁸

Expression of VPF/VEGF Receptors

As noted earlier, two high-affinity VPF/VEGF receptors have been described in vascular endothelium, flt-1 and KDR; both are transmembrane proteins with cytoplasmic tyrosine kinase domains.⁶⁹⁻⁷⁴ It is presumed, but not yet proven, that all of the effects of VPF/VEGF on vascular endothelium are mediated through one or both of these receptors.

In situ hybridization detects low level expression of both flt-1 and KDR mRNAs in the glomerular ECs of normal adult kidneys,⁶⁰ but signal is not detectable in most other normal adult tissues. However, at sites of VPF/VEGF overexpression, both flt-1 and KDR are commonly and strikingly overexpressed in adjacent microvascular ECs. Examples include various human and animal tumors,⁴¹⁻⁴⁴ the dermis in psoriasis⁴⁶ and contact dermatitis,⁴⁷ and the pannus of rheumatoid arthritis.^{38,50} Rarely, one or another VPF/VEGF receptor has been found to be expressed by cells other than ECs where their function, if any, is unclear.^{75,76}

Deposition of VPF/VEGF in Microvessels Adjacent to Sites of Its Overproduction

In addition to its detection in the cytoplasm of tumor cells, strong immunohistochemical staining for VPF/VEGF has been consistently demonstrated in adjacent hyperpermeable and replicating microvessels.^{41-43,77} However, these microvessels do not themselves express detectable VPF/VEGF mRNA by *in situ* hybridization, and therefore the observed immunohistochemical staining is thought to represent binding of VPF/VEGF secreted by adjacent tumor cells. Similar microvessel staining has been observed at non-neoplastic sites of VPF/VEGF overproduction and angiogenesis; eg, in the microvessels supplying rheumatoid arthritis pannus⁵⁰ and ovarian corpora lutea.⁵⁷

Recently, ultrastructural immunocytochemistry has localized tumor microvessel-bound VPF/VEGF more precisely to the abluminal surface of the EC plasma membrane and to VVOs.³⁶ This pattern of VPF/VEGF deposition could reflect binding to specific VPF/VEGF receptors. However, the intensity of the staining, the potency of VPF/VEGF action at low (sub-pmol/L to low nmol/L) concentrations, and the relatively small numbers of flt-1 and KDR receptors on individual ECs⁷⁸ (M Detmar, unpublished data) argue in favor of other possibilities, eg, binding to cell-surface proteoglycans in a manner analogous to that of basic fibroblast growth factor⁷⁹ or TGF- β .⁸⁰ Binding of VPF/VEGF to ECs could have functions other than cell signaling; eg, it may serve to keep VPF/VEGF localized, pre-

venting the widespread dissemination of a potent and potentially deleterious biological activity. In any event, VPF/VEGF immunostaining serves to identify blood vessels at sites of VPF/VEGF overproduction, distinguishing them from vessels elsewhere in the body. This finding has raised the possibility that VPF/VEGF may provide a useful target for tumor imaging and/or therapy.^{77,81}

Association Between Microvascular Hyperpermeability and Angiogenesis

The findings presented above link overexpression of VPF/VEGF (and its EC receptors) to angiogenesis as it occurs in many different pathological and physiological settings. This association makes intuitive sense in that VPF/VEGF is an angiogenic factor and as such might be expected to stimulate ECs to divide, migrate, and express matrix-degrading proteases, all properties characteristic of ECs in newly forming blood vessels. However, in addition to its other properties, VPF/VEGF renders microvessels hyperpermeable to circulating macromolecules, acting with great potency. An obvious question therefore arises: Is vascular hyperpermeability a consistent feature of angiogenesis? In fact, although rarely commented upon, the answer to this question is a categorical "yes". Published data from a number of different laboratories indicate that, wherever angiogenesis has been investigated, newly formed microvessels are hyperpermeable.

The most extensive evidence for an association between microvascular hyperpermeability and angiogenesis has come from studies of solid animal tumors. That the blood vessels supplying such tumors might be hyperpermeable was suggested some two decades ago by Gullino,⁸² who found that the concentrations of plasma proteins and water in tumor stroma were substantially higher than in the stroma of corresponding normal tissues. Moreover, direct permeability measurements with a wide variety of intravenously injected macromolecular tracers have convincingly demonstrated the innate leakiness to plasma proteins of the microvessels supplying tumors.^{2,3,81,83-91} Studies of ascites tumors have provided additional strong support for the inherent leakiness of tumor vessels.^{4,34} In addition, they have demonstrated high concentrations of immunoreactive and bioactive VPF/VEGF in tumor ascites fluid^{4,34,92} and have shown that vascular hyperpermeability precedes and accompanies angiogenesis.^{34,93} In all of the tumors that we have studied, vascular leakage has occurred primarily by way of VVOs and could not

be attributed to passage of tracers through inter-EC junctions or to trivial explanations such as EC injury.^{35,36,88}

It is also clear from the work of many different investigators that microvessels are hyperpermeable in pathological angiogenesis that is not associated with neoplasia, eg, in wound healing,^{45,94} rheumatoid arthritis,⁵⁰ psoriasis,^{46,95,96} delayed-type hypersensitivity,⁹⁷ etc. Finally, vascular hyperpermeability is also associated with the angiogenesis associated with such physiological events as corpus luteum formation in the ovary.^{56,57,98-101}

Microvascular Hyperpermeability as an Early, Mechanistic Step in the Induction of Angiogenesis

The demonstrated association between microvascular hyperpermeability and angiogenesis again raises a question: Does microvascular hyperpermeability play a mechanistic role in angiogenesis? Once again the answer is affirmative. We hypothesized more than a decade ago that plasma proteins extravasate from leaky blood vessels that supply tumors and form a new, provisional extracellular matrix that permits and indeed favors the inward migration of ECs and fibroblasts.^{91,102} Acting in concert, migrating ECs form new blood vessels and fibroblasts synthesize and secrete the matrix proteins, proteoglycans, and glycosaminoglycans that make up mature tumor stroma.^{3,91,102-105} According to this line of thought the extracellular matrix present in normal adult tissues might be expected to limit or even prohibit mesenchymal cell migration; this is not an unreasonable premise in that EC migration would be inappropriate in a mature tissue in which new blood vessel formation is contraindicated. Therefore, if new blood vessels and supporting connective tissue are to develop in adult tissues, as, eg, in response to the increased metabolic needs of a growing tumor, native tissue stroma must first be replaced with a new matrix that supports EC and fibroblast migration.

A large body of evidence now supports this hypothesis. Plasma proteins leaking from hyperpermeable blood vessels that supply animal tumors do in fact provide a matrix that favors mesenchymal cell migration. Within a few hours of transplant, pre-existing microvessels become measurably hyperpermeable to macromolecules and tumor-supplying microvessels eventually become 4 to 10-fold more permeable than their normal tissue counterparts.^{2,86,88} Moreover, plasma fibrinogen that extravasates at tumor sites rapidly clots to form crosslinked fibrin.^{2,3,86,106-108} This last finding indi-

cates, moreover, that the plasma protein extravasation that tumors induce is relatively nondiscriminating in that, to form an extravascular clot, a variety of plasma clotting proteins are required and must extravasate along with fibrinogen; these include at least clotting factors V, VII, X, and XIII, and prothrombin.^{109,110} The fibrin gel that is deposited in tumor stroma is itself modulated by plasmin, which is generated locally from another leaked plasma protein, plasminogen, by the action of tumor-(and perhaps EC-) secreted plasminogen activators.^{2,21,86,91,102,111} Plasmin also activates matrix metalloproteases, which are capable of digesting other matrix elements. The extent of fibrin deposition and its persistence over time vary extensively among different tumors, as does the amount of mature stroma generated by different tumors. Apparently these differences depend on quantitative balances among vascular hyperpermeability, clotting, and fibrinolysis that are unique to individual tumors.⁸⁶ These balances are further complicated because, as noted earlier, one of the actions of VPF/VEGF on ECs is to stimulate the synthesis of proteins that affect both coagulation (tissue factor)³⁹ and fibrinolysis (uPA, tPA, and PAI-1).²¹

Supportive evidence for our hypothesis also comes from *in vitro* studies that have provided direct evidence that crosslinked fibrin of the type deposited in tumors provides a matrix that supports and favors cell migration.^{112,113} Furthermore, implantation of crosslinked fibrin in guinea pigs, in the absence of tumor cells, induces the progressive ingrowth of new blood vessels and fibroblasts, which together generate vascularized stroma of the type found in many tumors.^{3,91,102,103,114} Fibrin exerts its pro-angiogenic effect at least in part by providing a favorable surface for cell adhesion and migration, presumably via its arg-gly-asp (RGD) sequence. Though not yet investigated, it is very likely that other circulating RGD-containing plasma proteins also extravasate from leaky blood vessels at sites of tumor growth and there contribute to the generation of an extracellular matrix that favors angiogenesis and new stroma formation; candidate plasma proteins include fibronectin and vitronectin.

If the hypothesis we have proposed for tumor angiogenesis is valid, it might also be expected to apply to the angiogenesis that occurs in non-neoplastic settings. In fact, this seems to be the case, and we (and others) have previously called attention to the many similarities between tumor stroma generation and wound healing.^{91,115} Other examples in which overexpression of VPF/VEGF and its receptors, plasma protein extravasation, and deposition of extravascu-

lar fibrin accompany angiogenesis include corpus luteum formation, psoriasis, contact allergy, and rheumatoid arthritis.^{45–47,50,56,94,97,116–121} Together these findings suggest that angiogenesis, whenever it occurs in adult tissues in response to diverse stimuli, is mediated by the triggering of a common pathway, initiated in every instance by VPF/VEGF overexpression and heightened vascular permeability.

Summary and Conclusions

VPF/VEGF is a multifunctional cytokine that contributes to angiogenesis by both direct and indirect mechanisms. On the one hand, VPF/VEGF stimulates the ECs lining nearby microvessels to proliferate, to migrate, and to alter their pattern of gene expression. On the other hand, VPF/VEGF renders these same microvascular ECs hyperpermeable so that they spill plasma proteins into the extravascular space, leading to the clotting of extravasated fibrinogen with deposition of a fibrin gel. Extravascular fibrin serves as a provisional matrix that favors and supports the ingrowth of new blood vessels and other mesenchymal cells that generate mature, vascularized stroma. These same principles apply in tumors, in several examples of non-neoplastic pathology, and in physiological processes that involve angiogenesis and new stroma generation. In all of these examples, microvascular hyperpermeability and the introduction of a provisional, plasma-derived matrix precede and accompany the onset of EC division and new blood vessel formation. It would seem, therefore, that tumors have “borrowed” fundamental mechanisms that developed in multicellular organisms for purposes of tissue defense, renewal, and repair.

VPF/VEGF, therefore, has taught us something new about angiogenesis; namely, that vascular hyperpermeability and consequent plasma protein extravasation are important, perhaps essential, elements in its generation. However, this finding raises a paradox. While VPF/VEGF induces vascular hyperpermeability, other potent angiogenic factors apparently do not, at least in subtoxic concentrations that are more than sufficient to induce angiogenesis.¹¹ Nonetheless, wherever angiogenesis has been studied, the newly generated vessels have been found to be hyperpermeable. How, therefore, do angiogenic factors other than VPF/VEGF lead to the formation of new and leaky blood vessels? We do not as yet have a complete answer to this question. One possibility is that at least some angiogenic factors mediate their effect by inducing or stimulating the expression of VPF/VEGF. In fact, there is already one clear example of this. TGF- α

is a potent angiogenic factor but does not itself increase microvascular permeability. However, TGF- α strikingly upregulates VPF/VEGF expression in cultured keratinocytes and is thought to be responsible, at least in part, for the overexpression of VPF/VEGF in psoriasis.⁴⁶ Moreover, overexpression of TGF- α , along with that of the EGF receptor with which it interacts, is characteristic of many malignant tumors, raising the possibility that TGF- α acts to stimulate VPF/VEGF expression in other types of epithelial cells and in this manner induces angiogenesis. Further studies that elucidate the crosstalk among various angiogenic factors are likely to contribute importantly to a better understanding of the mechanisms by which new blood vessels are formed in health and in disease.

Acknowledgments

The authors thank Dr. Donald R. Senger for helpful comments on the manuscript.

References

1. Senger D, Van De Water L, Brown L, Nagy J, Yeo K-T, Yeo T-K, Berse B, Jackman R, Dvorak A, Dvorak H: Vascular permeability factor (VPF, VEGF) in tumor biology. *Cancer Metastasis Rev* 1993, 12:303–324
2. Dvorak HF, Orenstein NS, Carvalho AC, Churchill WH, Dvorak AM, Galli SJ, Feder J, Bitzer AM, Rypysc J, Giovenco P: Induction of a fibrin-gel investment: an early event in line 10 hepatocarcinoma growth mediated by tumor-secreted products. *J Immunol* 1979, 122:166–174
3. Dvorak HF, Dvorak AM, Manseau EJ, Wiberg L, Churchill WH: Fibrin gel investment associated with line 1 and line 10 solid tumor growth, angiogenesis, and fibroplasia in guinea pigs. Role of cellular immunity, myofibroblasts, microvascular damage, and infarction in line 1 tumor regression. *J Natl Cancer Inst* 1979, 62:1459–1472
4. Senger DR, Galli SJ, Dvorak AM, Perruzzi CA, Harvey VS, Dvorak HF: Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* 1983, 219:983–985
5. Senger DR, Perruzzi CA, Feder J, Dvorak HF: A highly conserved vascular permeability factor secreted by a variety of human and rodent tumor cell lines. *Cancer Res* 1986, 46:5629–5632
6. Senger DR, Connolly D, Perruzzi CA, Alsup D, Nelson R, Leimgruber R, Feder J, Dvorak HF: Purification of a vascular permeability factor (VPF) from tumor cell conditioned medium. *Fed Proc* 1987, 46:2102
7. Connolly DT, Olander JV, Heuvelman D, Nelson R, Monsell R, Siegel N, Haymore BL, Leimgruber R,

- Feder J: Human vascular permeability factor. Isolation from U937 cells. *J Biol Chem* 1989, 264:20017-20024
8. Senger DR, Connolly DT, Van De Water L, Feder J, Dvorak HF: Purification and NH₂-terminal amino acid sequence of guinea pig tumor-secreted vascular permeability factor. *Cancer Res* 1990, 50:1774-1778
9. Brock TA, Dvorak HF, Senger DR: Tumor-secreted vascular permeability factor increases cytosolic Ca²⁺ and von Willebrand factor release in human endothelial cells. *Am J Pathol* 1991, 138:213-221
10. Senger DR, Dvorak HF: Vascular permeability factor: a protein mediator that induces tumor vessel hyperpermeability and promotes endothelial cell growth. *Human Astrocytomas*. Edited by PM Black and L Lampson. London, Blackwell Scientific Publications, 1993, pp 250-260
11. Connolly DT, Heuvelman DM, Nelson R, Olander JV, Eppley BL, Delfino JJ, Siegel NR, Leimgruber RM, Feder J: Tumor vascular permeability factor stimulates endothelial cell growth and angiogenesis. *J Clin Invest* 1989, 84:1470-1478
12. Ferrara N, Henzel WJ: Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells. *Biochem Biophys Res Commun* 1989, 161:851-858
13. Ferrara N, Houck K, Jakeman L, Winer J, Leung D: The vascular endothelial growth factor family of polypeptides. *J Cell Biochem* 1991, 47:211-218
14. Ferrara N, Houck K, Jakeman L, Leung D: Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocr Rev* 1992, 13:18-32
15. Gospodarowicz D, Abraham JA, Schilling J: Isolation and characterization of a vascular endothelial cell mitogen produced by pituitary-derived folliculo-stellate cells. *Proc Natl Acad Sci USA* 1989, 86:7311-7315
16. Houck KA, Ferrara N, Winer J, Cachianes G, Li B, Leung DW: The vascular endothelial growth factor family: identification of a fourth molecular species and characterization of alternative splicing of RNA. *Mol Endocrinol* 1991, 5:1806-1814
17. Plouet J, Schilling J, Gospodarowicz D: Isolation and characterization of a newly identified endothelial cell mitogen produced by AtT-20 cells. *EMBO J* 1989, 8:3801-3806
18. Sioussat TM, Dvorak HF, Brock TA, Senger DR: Inhibition of vascular permeability factor (vascular endothelial growth factor) with anti-peptide antibodies. *Arch Biochem Biophys* 1993, 301:15-20
19. Tischer E, Gospodarowicz D, Mitchell R, Silva M, Schilling J, Lau K, Crisp T, Fiddes JC, Abraham JA: Vascular endothelial growth factor. *Biochem Biophys Res Commun* 1989, 165:1198-1206
20. Favard C, Moukadiri H, Dorey C, Praloran V, Plouet J: Purification and biological properties of vasculotropin, a new angiogenic cytokine. *Biol Cell* 1991, 73:1-6
21. Pepper MS, Ferrara N, Orci L, Montesano R: Vascular endothelial growth factor (VEGF) induces plasminogen activators and plasminogen activator inhibitor-1 in microvascular endothelial cells. *Biochem Biophys Res Commun* 1991, 181:902-906
22. Unemori EN, Ferrara N, Bauer EA, Amento EP: Vascular endothelial growth factor induces interstitial collagenase expression in human endothelial cells. *J Cell Physiol* 1992, 153:557-562
23. Leung DW, Cachianes G, Kuang W-J, Goeddel DV, Ferrara N: Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 1989, 246:1306-1309
24. Wilting J, Christ B, Weich HA: The effects of growth factors on the day 13 chorioallantoic membrane (CAM): a study of VEGF₁₆₅ and PDGF-BB. *Anat and Embryol* 1992, 186:251-257
25. Folkman J, Shing Y: Angiogenesis. *J Biol Chem* 1992, 267:10931-10934
26. Conn G, Soderman DD, Schaeffer M-T, Wile M, Hatcher VB, Thomas KA: Purification of a glycoprotein vascular endothelial cell mitogen from a rat glioma-derived cell line. *Proc Natl Acad Sci USA* 1990, 87:1323-1327
27. Myoken Y, Kayada Y, Okamoto T, Kan M, Sato GH, Sato JD: Vascular endothelial cell growth factor (VEGF) produced by A-431 human epidermoid carcinoma cells and identification of VEGF membrane binding sites. *Proc Natl Acad Sci USA* 1991, 88:5819-5823
28. Yeo T-K, Senger DR, Dvorak HF, Freter L, Yeo K-T: Glycosylation is essential for efficient secretion but not for permeability-enhancing activity of vascular permeability factor (vascular endothelial growth factor). *Biochem Biophys Res Commun* 1991, 179:1568-1575
29. Maglione D, Guerriero V, Viglietto G, Delli-Bovi P, Persico MG: Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor. *Proc Natl Acad Sci USA* 1991, 88:9267-9271
30. Tischer E, Mitchell R, Hartman T, Silva M, Gospodarowicz D, Fiddes JC, Abraham JA: The human gene for vascular endothelial growth factor. Multiple protein forms are encoded through alternative exon splicing. *J Biol Chem* 1991, 266:11947-11954
31. Park J, Keller G-A, Ferrara N: The vascular endothelial growth factor (VEGF) isoforms: differential deposition into the subepithelial extracellular matrix and bioactivity of extracellular matrix-bound VEGF. *Mol Biol Cell* 1993, 4:1317-1326
32. Houck KA, Leung DW, Rowland AM, Winer J, Ferrara N: Dual regulation of vascular endothelial growth factor bioavailability by genetic and proteolytic mechanisms. *J Biol Chem* 1992, 267:26031-26037
33. Collins PD, Connolly DT, Williams TJ: Characterization of the increase in vascular permeability induced

- by vascular permeability factor *in vivo*. *Br J Pharmacol* 1993, 109:195–199
34. Nagy JA, Masse EM, Herzberg KT, Meyers MS, Yeo K-T, Yeo T-K, Sioussat TM, Dvorak HF: Pathogenesis of ascites tumor growth. Vascular permeability factor, vascular hyperpermeability and ascites fluid accumulation. *Cancer Res* 1995, 55:360–368
35. Kohn S, Nagy JA, Dvorak HF, Dvorak AM: Pathways of macromolecular tracer transport across venules and small veins. Structural basis for the hyperpermeability of tumor blood vessels. *Lab Invest* 1992, 67: 596–607
36. Qu-Hong, Nagy JA, Senger DR, Dvorak HF, Dvorak AM: Ultrastructural localization of vascular permeability factor/vascular endothelial growth factor (VPF/VEGF) to the abluminal plasma membrane and vesiculo-vacuolar organelles of tumor microvascular endothelium. *J Histochem Cytochem* 1995 (in press)
37. Rippe B, Haraldsson B: Transport of macromolecules across microvascular walls: the two-pore theory. *Physiol Rev* 1994, 74:163–219
38. Koch A, Harlow L, Haines G, Amento E, Unemori E, Wong W, Pope R, Ferrara N: Vascular endothelial growth factor. A cytokine modulating endothelial function in rheumatoid arthritis. *J Immunol* 1994, 152: 4149–4156
39. Clauss M, Gerlach M, Gerlach H, Brett J, Wang F, Familletti PC, Pan Y-CE, Olander JV, Connolly DT, Stern D: Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration. *J Exp Med* 1990, 172:1535–1545
40. Midy V, Plouet J: Vasculotropin/Vascular endothelial growth factor induces differentiation in cultured osteoblasts. *Biochem Biophys Res Commun* 1994, 199: 380–386
41. Brown LF, Berse B, Jackman RW, Tognazzi K, Manseau EJ, Senger DR, Dvorak HF: Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in adenocarcinomas of the gastrointestinal tract. *Cancer Res* 1993, 53: 4727–4735
42. Brown LF, Berse B, Jackman RW, Tognazzi K, Manseau EJ, Dvorak HF, Senger DR: Increased expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in kidney and bladder carcinomas. *Am J Pathol* 1993, 143: 1255–1262
43. Plate KH, Breier G, Weich HA, Risau W: Vascular endothelial growth factor is a potential tumour angiogenesis factor in human gliomas *in vivo*. *Nature* 1992, 359:845–848
44. Brown L, Berse B, Jackman R, Tognazzi K, Guidi A, Dvorak H, Senger D, Connolly J, Schnitt S: Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in breast cancer. *Hum Pathol* 1995, 26:86–91
45. Brown LF, Yeo K-T, Berse B, Yeo T-K, Senger DR, Dvorak HF, Van De Water L: Expression of vascular permeability factor (vascular endothelial growth factor) by epidermal keratinocytes during wound healing. *J Exp Med* 1992, 176:1375–1379
46. Detmar M, Brown L, Claffey K, Yeo K-T, Kocher O, Jackman R, Berse B, Dvorak H: Overexpression of vascular permeability factor and its receptors in psoriasis. *J Exp Med* 1994, 180:1141–1146
47. Brown LF, Olbricht SM, Berswe B, Jackman RW, Matsueda G, Tognazzi KA, Manseau EJ, Dvorak HF, Van De Water L: Overexpression of vascular permeability factor (VPF/VEGF) and its endothelial cell receptors in delayed hypersensitivity skin reactions. *J Immunol* 1995, 154:2801–2807
48. Ladoux A, Frelin C: Hypoxia is a strong inducer of vascular endothelial growth factor mRNA expression in the heart. *Biochem Biophys Res Commun* 1993, 195:1005–1010
49. Sharma H, Wunsch M, Schmidt M, Schott R, Kandolf R, Schaper W: Expression of angiogenic growth factors in the collateralized swine myocardium. *EXS* 1992, 61:255–260
50. Fava R, Olsen N, Spencer-Green G, Yeo K-T, Yeo T-K, Berse B, Jackman R, Senger D, Dvorak H, Brown L: Vascular permeability factor/endothelial growth factor (VPF/VEGF): Accumulation and expression in human synovial fluids and rheumatoid synovial tissue. *J Exp Med* 1994, 180:341–346
51. Sharkey A, Charnock-Jones D, Boocock C, Brown K, Smith S: Expression of mRNA for vascular endothelial growth factor in human placenta. *J Reprod Fertil* 1993, 99:609–615
52. Jackson MR, Carney EW, Lye SJ, Ritchie JW: Localization of two angiogenic growth factors (PDECGF and VEGF) in human placentae throughout gestation. *Placenta* 1994, 15:341–353
53. Shweiki D, Itin A, Neufeld G, Gitay GH, Keshet E: Patterns of expression of vascular endothelial growth factor (VEGF) and VEGF receptors in mice suggest a role in hormonally regulated angiogenesis. *J Clin Invest* 1993, 91:2235–2243
54. Breier G, Albrecht U, Sterrer S, Risau W: Expression of vascular endothelial growth factor during embryonic angiogenesis and endothelial cell differentiation. *Development* 1992, 114:521–532
55. Shifren JL, Doldi N, Ferrara N, Mesiano S, Jaffe RB: In the human fetus, vascular endothelial growth factor is expressed in epithelial cells and myocytes, but not vascular endothelium: Implications for mode of action. *J Clin Endocrinol Metab* 1994, 79:316–322
56. Cullinan BK, Koos RD: Vascular endothelial growth factor/vascular permeability factor expression in the rat uterus: rapid stimulation by estrogen correlates with estrogen-induced increases in uterine capillary permeability and growth. *Endocrinology* 1993, 133: 829–837
57. Kamat BR, Brown LF, Manseau EJ, Senger DR, Dvorak HF: Expression of vascular permeability

- factor/vascular endothelial growth factor by human granulosa and theca lutein cells: role in corpus luteum development. *Am J Pathol* 1995, 146:157–165
58. Phillips HS, Hains J, Leung DW, Ferrara N: Vascular endothelial growth factor is expressed in rat corpus luteum. *Endocrinology* 1990, 127:965–967
59. Berse B, Brown LF, Van De Water L, Dvorak HF, Senger DR: Vascular permeability factor (vascular endothelial growth factor) gene is expressed differentially in normal tissues, macrophages, and tumors. *Mol Biol Cell* 1992, 3:211–220
60. Brown LF, Berse B, Tognazzi K, Manseau EJ, Van De Water L, Senger DR, Dvorak HF, Rosen S: Vascular permeability factor mRNA and protein expression in human kidney. *Kidney Int* 1992, 42:1457–1461
61. Brown L, Yeo K-T, Berse B, Morgentaler A, Dvorak H, Rosen S: Vascular permeability factor (vascular endothelial growth factor) is strongly expressed in the normal male genital tract and is present in substantial quantities in semen. *J Urology* 1995 (in press)
62. Dolecki GJ, Connolly DT: Effects of a variety of cytokines and inducing agents on vascular permeability factor mRNA levels in U937 cells. *Biochem Biophys Res Commun* 1991, 180:572–578
63. Finkenzeller G, Marme D, Weich HA, Hug H: Platelet-derived growth factor-induced transcription of the vascular endothelial growth factor gene is mediated by protein kinase C. *Cancer Res* 1992, 52:4821–4823
64. Brogi E, Wu T, Namiki A, Isner JM: Indirect angiogenic cytokines upregulate VEGF and bFGF gene expression in vascular smooth muscle cells, whereas hypoxia upregulates VEGF expression only. *Circulation* 1994, 90:649–652
65. Pertovaara L, Kaipainen A, Mustonen T, Orpana A, Ferrara N, Saksela O, Alitalo K: Vascular endothelial growth factor is induced in response to transforming growth factor- β in fibroblastic and epithelial cells. *J Biol Chem* 1994, 269:6271–6274
66. Goldberg M, Schneider T: Similarities between the oxygen sensing mechanisms regulating the expression of vascular endothelial growth factor and erythropoietin. *J Biol Chem* 1994, 269:4355–4359
67. Shweiki D, Itin A, Soffer D, Keshet E: Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature* 1992, 359:843–845
68. Koos RD, Olson CE: Hypoxia stimulates expression of the gene for vascular endothelial growth factor (VEGF), a putative angiogenic factor, by granulosa cells of the ovarian follicle, a site of angiogenesis (abstract). *J Cell Biol* 1991, 115:421a
69. Terman BI, Carrion ME, Kovacs E, Rasmussen BA, Eddy RL, Shows TB: Identification of a new endothelial cell growth factor receptor tyrosine kinase. *Oncogene* 1991, 6:1677–1683
70. Terman BI, Dougher-Vermazen M, Carrion ME, Dimitrov D, Armellino DC, Gospodarowicz D, Böhlen P: Identification of the KDR tyrosine kinase as a receptor for vascular endothelial cell growth factor. *BBRC* 1992, 187:1579–1586
71. de Vries C, Escobedo JA, Ueno H, Houck K, Ferrara N, Williams LT: The *fms*-like tyrosine kinase, a receptor for vascular endothelial growth factor. *Science* 1992, 255:989–991
72. Matthews W, Jordan C, Gavin M, Jenkins N, Copeland N, Lemischka I: A receptor tyrosine kinase cDNA isolated from a population of enriched primitive hematopoietic cells and exhibiting close genetic linkage to *c-kit*. *Proc Natl Acad Sci USA* 1991, 88:9026–9030
73. Millauer B, Wizigmann-Voos S, Schnurch H, Martinez R, Moller N, Risau W, Ullrich A: High affinity VEGF binding and developmental expression suggest Flk-1 as a major regulator of vasculogenesis and angiogenesis. *Cell* 1993, 72:835–846
74. Quinn T, Peters K, DeVries C, Ferrara N, Williams L: Fetal liver kinase 1 is a receptor for vascular endothelial growth factor and is selectively expressed in vascular endothelium. *Proc Natl Acad Sci USA* 1993, 90:7533–7537
75. Charnock-Jones D, Sharkey A, Boock C, Ahmed A, Plevin R, Ferrara N, Smith S: Vascular endothelial growth factor receptor localization and activation in human trophoblast and choriocarcinoma cells. *Biol Reprod* 1994, 51:524–530
76. Gitay-Goren H, Halaban R, Neufeld G: Human melanoma cells but not normal melanocytes express vascular endothelial growth factor receptors. *BBRC* 1993, 190:702–709
77. Dvorak HF, Sioussat TM, Brown LF, Berse B, Nagy JA, Sotrel A, Manseau EJ, Van De Water L, Senger DR: Distribution of vascular permeability factor (vascular endothelial growth factor) in tumors: concentration in tumor blood vessels. *J Exp Med* 1991, 174:1275–1278
78. Vaisman N, Gospodarowicz D, Neufeld G: Characterization of the receptors for vascular endothelial growth factor. *J Biol Chem* 1990, 265:19461–19466
79. Yayon A, Klagsbrun M, Esko JD, Leder P, Ornitz DM: Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell* 1991, 64:841–848
80. Andres JL, Stanley K, Chieftetz S, Massagué J: Membrane-anchored and soluble forms of betaglycan, a polymorphic proteoglycan that binds transforming growth factor- β . *J Cell Biol* 1989, 109:3137–3145
81. Dvorak HF, Nagy JA, Dvorak AM: Structure of solid tumors and their vasculature: implications for therapy with monoclonal antibodies. *Cancer Cells* 1991, 3:77–85
82. Gullino PM: Extracellular compartments of solid tumors, *Cancer: A Comprehensive Treatise*. Edited by

- FF Becker. New York, Plenum Press, 1975, pp 327–354
83. Underwood JC, Carr I: The ultrastructure and permeability characteristics of the blood vessels of a transplantable rat sarcoma. *J Pathol* 1972, 107:157–166
84. Peterson H-I: Tumor Blood Circulation: Angiogenesis, Vascular Morphology and Blood Flow of Experimental and Human Tumors. Boca Raton, CRC Press, Inc., 1979, pp 1–229
85. O'Connor SW, Bale WF: Accessibility of circulating immunoglobulin G to the extravascular compartment of solid rat tumors. *Cancer Res* 1984, 44:3719–3723
86. Dvorak HF, Harvey VS, McDonagh J: Quantitation of fibrinogen influx and fibrin deposition and turnover in line 1 and line 10 guinea pig carcinomas. *Cancer Res* 1984, 44:3348–3354
87. Dvorak HF, Nagy JA, Berse B, Brown LF, Yeo K-T, Yeo T-K, Dvorak AM, Van De Water L, Sioussat TM, Senger DR: Vascular permeability factor, fibrin, and the pathogenesis of tumor stroma formation. *Ann NY Acad Sci* 1992, 667:101–111
88. Dvorak HF, Nagy JA, Dvorak JT, Dvorak AM: Identification and characterization of the blood vessels of solid tumors that are leaky to circulating macromolecules. *Am J Pathol* 1988, 133:95–109
89. Gerlowski LE, Jain RK: Microvascular permeability of normal and neoplastic tissues. *Microvasc Res* 1986, 31:288–305
90. Heuser LS, Miller FN: Differential macromolecular leakage from the vasculature of tumors. *Cancer* 1986, 57:461–464
91. Dvorak HF: Tumors: wounds that do not heal. Similarities between tumor stroma generation and wound healing. *N Engl J Med* 1986, 315:1650–1659
92. Yeo K-T, Wang HH, Nagy JA, Sioussat TM, Ledbetter SR, Hoogewerf AJ, Zhou Y, Masse EM, Senger DR, Dvorak HF, Yeo T-K: Vascular permeability factor (vascular endothelial growth factor) in guinea pig and human tumor and inflammatory effusions. *Cancer Res* 1993, 53:2912–2918
93. Nagy J, Morgan E, Herzberg K, Manseau E, Dvorak A, Dvorak H: Pathogenesis of ascites tumor growth. Angiogenesis, vascular remodeling and stroma formation in the peritoneal lining. *Cancer Res* 1995, 55:376–385
94. Breuing K, Eriksson E, Liu P, Miller DR: Healing of partial thickness porcine skin wounds in a liquid environment. *J Surg Res* 1992, 52:50–58
95. Braverman IM, Keh YA: Three-dimensional reconstruction of endothelial cell gaps in psoriatic vessels and their morphologic identity with gaps produced by the intradermal injection of histamine. *J Invest Dermatol* 1986, 86:577–581
96. Bull RH, Bates DO, Mortimer PS: Intravital video-capillaroscopy for the study of the microcirculation in psoriasis. *Br J Dermatol* 1992, 126:436–445
97. Colvin R, Dvorak H: Role of the clotting system in cell-mediated hypersensitivity. II. Kinetics of fibrinogen/fibrin accumulation and vascular permeability changes in tuberculin and cutaneous basophil hypersensitivity reactions. *J Immunology* 1975, 114:377–387
98. Morris B, Sass M: The formation of lymph in the ovary. *Proc R Soc Lond B Biol* 1966, 164:577–591
99. Moor RM, Seamark RF: Cell signaling, permeability, and microvasculatory changes during antral follicle development in mammals. *J Dairy Sci* 1986, 69:927–943
100. Okuda Y, Okamura H, Kanzaki H, Takenaka A, Morimoto K, Nishimura T: An ultrastructural study of capillary permeability of rabbit ovarian follicles during ovulation using carbon tracer. *Acta Obstet Gynecol Jpn* 1980, 32:859–867
101. Payer A: Permeability of ovarian follicles and capillaries in mice. *Am J Anat* 1975, 142:295–301
102. Dvorak HF, Senger DR, Dvorak AM: Fibrin as a component of the tumor stroma: origins and biological significance. *Cancer Metastasis Rev* 1983, 2:41–73
103. Dvorak HF, Form DM, Manseau EJ, Smith BD: Pathogenesis of desmoplasia. I. Immunofluorescence identification and localization of some structural proteins of line 1 and line 10 guinea pig tumors and of healing wounds. *J Natl Cancer Inst* 1984, 73:1195–1205
104. Yeo T-K, Brown L, Dvorak HF: Alterations in proteoglycan synthesis common to healing wounds and tumors. *Am J Pathol* 1991, 138:1437–1450
105. Yeo T-K, Dvorak HF: Tumor stroma. *Diagnostic Immunopathology*. Edited by R Colvin, A Bhan, and R McCluskey. New York, Raven Press, 1995, pp 685–697
106. Dvorak HF, Senger DR, Dvorak AM, Harvey VS, McDonagh J: Regulation of extravascular coagulation by microvascular permeability. *Science* 1985, 227:1059–1061
107. Brown LF, Asch B, Harvey VS, Buchinski B, Dvorak HF: Fibrinogen influx and accumulation of cross-linked fibrin in mouse carcinomas. *Cancer Res* 1988, 48:1920–1925
108. Brown LF, Van De Water L, Harvey VS, Dvorak HF: Fibrinogen influx and accumulation of cross-linked fibrin in healing wounds and in tumor stroma. *Am J Pathol* 1988, 130:455–465
109. Dvorak HF, Quay SC, Orenstein NS, Dvorak AM, Hahn P, Bitzer AM, Carvalho AC: Tumor shedding and coagulation. *Science* 1981, 212:923–924
110. Dvorak HF, Van De Water L, Bitzer AM, Dvorak AM, Anderson D, Harvey VS, Bach R, Davis GL, De Wolf W, Carvalho ACA: Procoagulant activity associated with plasma membrane vesicles shed by cultured tumor cells. *Cancer Res* 1983, 43:4434–4442
111. Danø K, Andreassen PA, Grøndahl-Hansen J, Kristensen P, Nielsen LS, Skriver L: Plasminogen activators, tissue degradation, and cancer. *Adv Cancer Res* 1985, 44:139–266
112. Lanir N, Ciano PS, Van De Water L, McDonagh J,

- Dvorak AM, Dvorak HF: Macrophage migration in fibrin gel matrices. II. Effects of clotting factor XIII, fibronectin, and glycosaminoglycan content on cell migration. *J Immunol* 1988, 140:2340–2349
113. Brown LF, Lanir N, McDonagh J, Czarnecki K, Estrella P, Dvorak AM, Dvorak HF: Fibroblast migration in fibrin gel matrices. *Am J Pathol* 1993, 142:273–283
114. Dvorak HF, Harvey VS, Estrella P, Brown LF, McDonagh J, Dvorak AM: Fibrin containing gels induce angiogenesis. Implications for tumor stroma generation and wound healing. *Lab Invest* 1987, 57:673–686
115. Haddow A: Molecular repair, wound healing, and carcinogenesis: tumor production a possible over-healing? *Adv Cancer Res* 1972, 16:181–234
116. Braverman IM, Yen A: Microcirculation in psoriatic skin. *J Invest Dermatol* 1974, 62:493–502
117. Brody I: Dermal and epidermal involvement in the evolution of acute eruptive guttate psoriasis vulgaris. *J Invest Dermatol* 1984, 82:465–470
118. Parent D, Bernard BA, Desbas C, Heenen M, Darmon MY: Spreading of psoriatic plaques: alteration of epidermal differentiation precedes capillary leakiness and anomalies in vascular morphology [see comments]. *J Invest Dermatol* 1990, 95:333–340
119. Salo OP, Kousa M, Mustakallio KK, Lassus A: Demonstration of fibrin in skin diseases. II. Psoriasis. *Acta Derm Venereol (Stockh)* 1972, 52:295–297
120. Staberg B, Worm AM, Klemp P, Rossing N: Transvascular transport and distribution of fluid and protein in psoriasis. *J Am Acad Dermatol* 1983, 8:193–199
121. Worm AM, Rossing N: Microvascular protein leakage in extensive skin diseases: aspects of the transport mechanisms. *J Invest Dermatol* 1980, 75:302–305